



## Correlation of HbA1c with serum LDL in newly diagnosed type 2 DM patients

Darshna Jain<sup>1</sup>, Prachi Paliwal<sup>2</sup>, Manoj Narayan Paliwal<sup>3</sup>

<sup>1</sup> Assistant Professor, Biochemistry, Govt. Medical College, Ratlam, Madhya Pradesh, India

<sup>2</sup> Associate Professor, Biochemistry, SAIMS, Indore, Madhya Pradesh, India

<sup>3</sup> Professor, Biochemistry, Govt. Medical College, Ratlam, Madhya Pradesh, India

\*Corresponding Author: Prachi Paliwal

### Abstract

**Background:** Glycated hemoglobin (HbA1c) is considered as a gold-standard measure of chronic glycemia in diabetic patients. HbA1c is a main indicator for mean blood glucose level. Dyslipidemia especially high LDL, is common in diabetes mellitus and is strongly associated with poor glycemic control.

**Method:** A retrospective study was carried out to determine the LDL levels in newly diagnosed type 2 diabetics. A total of 100 newly diagnosed type 2 diabetics were enrolled in our study. Relevant patient data was collected from hospital information system, SAIMS medical college, Indore.

**Result:** In our study, maximum patients (53.00%) were from 46-60 years, followed by, 25.00% patients were more than 60 years and 23.00% patients were less than 45 years of age. 45% patients were female and 55% patients were male. The association between HbA1c and LDL was found statistically significant.

**Conclusion:** It was concluded from the results of this study that type 2 diabetic patients are more prone to dyslipidemia. In newly diagnosed type 2 diabetes mellitus, high LDL levels were found in patients with high HbA1C levels.

**Keywords:** type 2 diabetes mellitus, HbA1c, LDL

### Introduction

One of the important cardiovascular risk factors in type 2 diabetes is dyslipidemia. The composition of lipids in diabetic dyslipidemia is more atherogenic than in dyslipidemia in general [1, 2]. The term diabetic dyslipidemia comprises a triad of raised triglycerides, reduced high density lipoprotein (HDL) and excess of small, dense low density lipoprotein (LDL) [3]. Every one of these dyslipidemic features are associated with an increased risk of cardiovascular disease (CVD). Increased hepatic secretion of large triglyceride-rich VLDL and impaired clearance of VLDL is central to the pathophysiology of this dyslipidemia [4]. The contribution of LDL to CVD risk has been much debated in the past, with many important prospective studies observing an association between elevated LDL levels and CVD risk [5]. This independent association with long term all-cause mortality supports the idea that serum LDL could play a role in type 2 diabetic patients mortality risk [2].

In the present study, we have aimed to study the LDL abnormalities in newly diagnosed type 2 diabetics. Such an assessment will enable earlier detection and treatment of these LDL derangements thereby minimizing the cardiovascular morbidity and mortality that can occur.

### Methodology

**Type of Study:** A retrospective study

**Sampling Methods:** Random sampling

### Inclusion Criteria

All patients of age >30years who have been diagnosed as new cases of type 2 diabetes in the OPD and IPD within the last 3 months [using the ADA (American Diabetes Association) criteria] were included.

### Exclusion Criteria

- Patients on steroids
- Type 1 diabetics
- Patients on antipsychotic medications
- Known cases of active hypothyroidism
- Known cases of Cushing's syndrome were excluded from the study.
- Unavailability of data on LDL and HbA1C levels.

### Data Collection

A retrospective study was carried out to determine the LDL levels in newly diagnosed type 2 diabetics. A total of 100 newly diagnosed type 2 diabetics from the OPD and IPD, whose LDL and HbA1C levels were available, were enrolled in our study. Based on the HbA1c levels, patients were divided into 3 categories such as, HbA1C<7, HbA1C 7-8 and HbA1C <8. Fasting lipid profile levels were measured in these patients

### Statistical Analysis

Continuous variables were expressed using mean, standard deviation, range and mean while categorical variables were expressed in terms of percentages. Test of significance was done using student t test for normally distributed continuous variables. Mann Whitney test was done for not normally distributed continuous variables. P value less than 0.05 was considered as statistically significant. All the statistical analysis were done using the software SPSS 23.0 (trial version).

## Results

**Table 1:** Socio-demographic profile

Age group in yrs	No of cases	Percentage
<45	27	27.00
46-60	53	53.00
>60	20	25.00
Male: female	55:45	

Mean age=53.27±10.97 years

In our study, maximum patients (53.00%) were from 46-60 years, followed by, 25.00% patients were more than 60 years and 23.00% patients were less than 45 years of age. 45% patients were female and 55% patients were male.

**Table 2:** Association between HbA1c and LDL level

Hb1Ac	LDL (mg/dl)		p-value
	Mean	SD	
<7 %	122.57	18.6	0.001
7-8 %	150.54	18.94	0.001
>8 %	168.8	13.02	0.001

The association between HbA1c and LDL was found statistically significant.

## Discussion

In our study, maximum patients (53.75%) were from 46-60 years followed by, 25% patients were more than 60 years and 21.25% patients were less than 45 years of age. This shows type 2 diabetes begins typically in middle life or later, the prevalence rises with age. This is consistent with studies published by ADA (2010). This also implies that the impact of age as a risk factor for diabetes cannot be overemphasized as this trend has been demonstrated in most study populations around the world [6].

The present study shows 55% male participants and 45% female participants. This differs from study done by Mohammad Al Shafee [7] in Oman in which, out of 1313 participants, 490 were male and 823 were female participants. So, we assume that, in Indore, firstly men have more access to outside culture and tend to seek more healthcare attention than female and secondly men have more participation owing to decreased female sex ratio.

In our study, the association between HbA1c and LDL was found statistically significant. In a study by H. Surekha Rani Et.al., it is observed that FBS and PPBS, Cholesterol, VLDL, LDLs, TGs were high and the levels of HDLs were low compared to controls [2]. Studies done by Maharjan *et al* [8], Cohen *et al* [9], and Arab *et al* [10], reported significant correlations between glycosylated hemoglobin and TG, TC, LDL and FBS and non-significant correlation with HDL.

Lipid abnormalities are common in diabetic patients and frequently seen in patients with type-2 diabetic mellitus. The abnormal lipid profile observed in type 2 Diabetes mellitus is said to be related to insulin resistance, which leads to increased release of free fatty acids from fatty tissue, impaired insulin dependent muscle uptake of free fatty acids and increase fatty acid release to the hepatic tissue [11]. It has been closely associated with diabetic dyslipidemia, hypertension and enormous risk to cardiovascular diseases [12]. Chronic hyperglycemia causes glycation of apolipoproteins and interferes with the normal pathways of lipoprotein metabolism [13]. Dyslipidemia which includes elevated levels of LDL and triacylglycerol and low level of HDL are more common among the diabetics.. In the present study, the results showed that the lipid profile were higher in

diabetic patients and that they were in agreement with the finding of Wexler *et al.*, 2005 [14]. Study by Elizabeth et.al, observed that LDL and HDL cholesterol were significantly associated with HbA1c [15]. In the present study we found significantly increased levels of LDL with higher HbA1c values.

Small sample size and the cause-effect relationship between the LDL and HbA1C levels are the limiting factors in our study. Here, only the association between the LDL and HbA1C levels could be established. So, studies with larger sample size and prospective aspect are needed for further confirmation.

## Conclusion

It was concluded from the results of this study that type 2 diabetic patients are more prone to dyslipidemia. In newly diagnosed type 2 diabetes mellitus, high LDL levels were found in patients with high HbA1C levels.

## Reference

1. Lt Gen SR Mehta, VSM, Col AS Kashyap, Lt Col S Das. Diabetes Mellitus in India: The Modern Scourge. MJAFI,2009;65:50-4.
2. Surekha Rani H, Madhavi G, Ramachandra Rao V, Sahay BK, Jyothy A. Risk Factors for Coronary Heart Disease in Type II DM. Indian Journal of Clinical Biochemistry,2005;20(2):75-80.
3. Zachary T, Bloomgarden. Cardiovascular Disease, Neuropathy and Retinopathy. Diabetes care,2009;32:64-68.
4. Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR *et al.* Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: The Atherosclerosis Mahantesh Patil et. al. Int J Cur Res Rev, March 2014/ Vol 06 (06) Page 24 Risk in Communities (ARIC) Study, 1987– 1993. Am J Epidemiol,1997;146:483-94.
5. Burtis Carl A, Ashwood Edward R, Bruns David E. Tietz Textbook of clinical chemistry and molecular diagnostics. 4th Ed. New Delhi: Elsevier publishers; 2008. Chapter 26, Lipids, lipoproteins, apolipoproteins and other cardiovascular risk factors,2008;26:903-83.
6. American Diabetes Association (2010) Position Statement; Standards of Medical Care in Diabetes-2010. Diabetes Care,2010;33(1):11-61.
7. Mohammed A Al-Shafee, Kamlesh Bhargava, Yahya Mohammed Al-Farsi, Shirley Mcilvenny. Prevalence of pre-diabetes and associated risk factors in adult Omani population Int J Diab Dev Ctries,2011;31:166-174.
8. Maharjan P, Pandeya D, Joshi G, Hona S, Bhatta B, Hamza A *et al.* Glycated Hemoglobin (Hba1C) Is a Predictor of Dyslipidemia in Type 2 Diabetes Nepalese Patients. Int J Adv Res (Indore),2017;5:113-121.
9. Cohen J. Statistical power analysis for the behavioral sciences. Taylor & Francis, Routledge, 1988.
10. Arab GA, Zahedi M, Kazemi V, Sanagoo A, Azimi M. Correlation between Hemoglobin A1C and Serum Lipid Profile in Type 2 Diabetic Patients Referred to the Diabetes Clinic in Gorgan. Iran J Clin Basic Res,2018;2:26-31.
11. Boden G; Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. Diabetes, 1997;46(1):4755.
12. Mgonda YM, Ramaiya KL, Swai AB, McLarty DG,

- Alberti KG. Insulin resistance and hypertension in non-obese Africans in Tanzania. *Hypertension*, 1998;31(1):114-118.
13. Carmena R. High risk of lipoprotein dysfunction in type 2 diabetes mellitus. *Rev EspCardiolSupl*,2008;8:18-24.
  14. Wexler DJ, Grant RW, Meigs JB, Nathan DM, Cagliero, E. Sex disparities in the treatment of cardiac risk factors in patients with type 2 diabetes. *Diabetes Care*,2005;28:514-520.
  15. Elizabeth Selvin, Josef Coresh, Sherita Golden H, Lori L Boland, Frederick Brancati L, Michael Steffes W. Glycemic Control, Atherosclerosis, and Risk Factors for Cardiovascular Disease in Individuals with Diabetes. *Diabetes Care*,2005;28(8):1965-73.